CONTENTS

| (Abstracts/contents lists published in: Chem Abstr.; Curr. Contents; Phys. Chem. Earth Sci.; Life Stexts are incorporated in CJELSEVIER, a file in the chemical Journals Online database which is available STN® International.) | |
|---|-----|
| Conformational analysis of isopropylidene-protected C-glycosyl derivatives of 3-deoxy-D-manno-2-octulosonic acid (Kdo) in the solid state and in solution | |
| M. Orbe, K. Luthman, T. Wåglund (Uppsala, Sweden), A. Claesson (Södertälje, Sweden), and I. Csöregh (Stockholm, Sweden) | 1 |
| Determination of the degree of N-acetylation and the distribution of N-acetyl groups in partially N-deacetylated chitins (chitosans) by high-field n.m.r. spectroscopy | |
| K. M. Varum, M. W. Anthonsen, H. Grasdalen, and O. Smidsrød (Trondheim, Norway) | 17 |
| Synthesis of two phosphate-containing "heptasaccharide" fragments of the capsular polysaccharides of Streptococcus pneumoniae types 6A and 6B | |
| T. M. Slaghek, A. A. M. Maas, J. P. Kamerling, and J. F. G. Vliegenthart (Utrecht, The | |
| Netherlands) | 25 |
| Analysis of positions of substitution of O-carboxymethyl groups in partially O-carboxymethylated cellulose by the reductive-cleavage method | |
| S. G. Zeller, G. W. Griesgraber, and G. R. Gray (Minneapolis, MN, U.S.A.) | 41 |
| Synthesis and mass spectra of 4-O-acetyl-1,5-anhydro-2,3,6-tri-O-(methoxycarbonylmethyl)-D-glucitol and the positional isomers of 4-O-acetyl-1,5-anhydro-di-O-(methoxycarbonylmethyl)-O-methyl-D-glucitol and 4-O-acetyl-1,5-anhydro-O-(methoxycarbonylmethyl)-di-O-methyl-D-glucitol | |
| S. G. Zeller, G. W. Griesgraber, and G. R. Gray (Minneapolis, MN, U.S.A.) | 47 |
| The design and synthesis of antibody binding site probes: three pentasaccharide analogues of the Brucella A antigen prepared by activation in situ of thioglycosides with bromine | |
| J. Kihlberg, E. Eichler, and D. R. Bundle (Ottawa, Ont., Canada) | 59 |
| Structural characterization of a new anticoagulant fucan sulfate from the brown seaweed Ecklonia kurome | |
| T. Nishino, T. Nagumo (Kanagawa, Japan), H. Kiyohara, and H. Yamada (Tokyo, Japan) | 77 |
| L-Guluronan-specific alginate lyase from a marine bacterium associated with Sargassum | |
| B. J. Brown and J. F. Preston, III (Gainesville, FL, U.S.A.) | 91 |
| Glucuronoxylomannan of Cryptococcus neoformans serotype B: structural analysis by gas-liquid | |
| chromatography—mass spectrometry and ¹³ C-nuclear magnetic resonance spectroscopy S. H. Turner and R. Cherniak (Atlanta, GA, U.S.A.) | 103 |
| A new undecasaccharide subunit of xyloglucans with two α-L-fucosyl residues | |
| M. Hisamatsu, G. Impallomeni, W. S. York, P. Albersheim, and A. G. Darvill (Athens, GA, U.S.A.) | 117 |
| Notes | |
| Votes The identification by ¹ H- and ¹³ C-n.m.r. spectroscopy of sucrose, 1-kestose, and neokestose in | |
| mixtures present in plant extracts | |
| A. De Bruyn (Gent, Belgium) and J. van Loo (Tienen, Belgium) | 131 |

| 3-Amino-5-C-phenyl-D-altrofuranose and 3-amino-5-C-[3-carboxy-4-(carboxymethyl)-2-oxo-3-cy-clohexen-1-yl]-D-altrofuranose, possible intermediates for synthesis of the anthracycline antibiotic decilorubicin | |
|---|-----|
| T. Izawa, Y. Nishimura, and S. Kondo (Tokyo, Japan) | 13 |
| Synthesis of pseudo-laminarabiose, -cellobiose, and -maltose (D-glucopyranosyl 5a-carba-D- and L-glucopyranoses) | |
| S. Ogawa, I. Sugawa, and Y. Shibata (Yokohama, Japan) | 14 |
| Glycosyl 1-piperidinecarbodithioates in the synthesis of glycosides P. Fügedi, P. J. Garegg, S. Oscarson, G. Rosen, and B. A. Silwanis (Stockholm, Sweden) | 15 |
| Observation of a stable orthoacyl fluoride in the reaction of a reducing sugar with diethylamino- sulfur trifluoride | |
| M. H. E. Griffith and O. Hindsgaul (Edmonton, Alta., Canada) | 163 |
| Autoxidative degradation of Amadori compounds in the presence of copper ion S. Kawakishi, J. Tsunehiro, and K. Uchida (Nagoya, Japan) | 167 |
| Preparation of acetylated 2,6-anhydrohept(hex)-2-enononitriles (1-cyano-2-hydroxyglycals) L. Somsák, E. Papp. G. Batta, and I. Farkas (Debrecen, Hungary) | 173 |
| Selective 3-O-allylation and 3-O-benzylation of methyl α -D-manno-, α -L-rhamno- and β -L-fuco-pyranoside | |
| G. Yang, F. Kong, and S. Zhou (Beijing, China) | 179 |
| Structural studies of the Escherichia coli 01A O-polysaccharide, using the computer program CASPER | |
| H. Baumann, PE. Jansson, L. Kenne, and G. Widmalm (Stockholm, Sweden) | 183 |
| Papers Structure and solution conformations of a cyclic trisaccharide from high-resolution n.m.r. spectroscopy and molecular modelling | |
| G. Bonas, M. R. Vignon (Grenoble, France), and S. Pérez (Nantes, France) | 191 |
| Ring-opening kinetics of the D-pentofuranuronic acids J. Wu and A. S. Serianni (Notre Dame, IN, U.S.A.) | 207 |
| Assignment of structures to oligosaccharides produced by enzymic degradation of a β -D-glucan from | |
| barley by ¹ H- and ¹³ C-n.m.r. spectroscopy K. Bock, J. Ø. Duus (Valby, Denmark), B. Norman, and S. Pedersen (Bagsvaerd, Denmark) | 219 |
| Mechanism of oxidation of reducing sugars by osmium tetraoxide in alkaline medium by the stopped-flow technique | |
| H. S. Singh, B. Singh, and A. K. Singh (Allahabad, India) | 235 |
| Branched-chain thiosugars: temperature effects in the acetolytic deacetalation of furanose sugars S. H. Kawai, J. Chin, and G. Just (Montreal, Que., Canada) | 245 |
| Synthesis of a spacer-containing repeating unit of the capsular polysaccharide of Streptococcus | |
| pneumoniae type 23F A. M. P. van Steijn, J. P. Kamerling, and J. F. G. Vliegenthart (Utrecht, The Netherlands) | 261 |
| A ¹³ C-n.m.r. study of the tungstate and molybdate complexes of perseitol, galacitol, and D-mannitol S. Chapelle and JF. Verchère (Mont-Saint-Aignan, France) | 279 |

| Synthesis of 3-(D-lyxofuranosyl)pyrazoles by trifluoracetic acid-catalysed cyclodehydration of 3-(D- galacto-pentitol-1-yl)pyrazoles | |
|--|-----|
| M. Gómez-Guillén and J. M. Lassaletta Simon (Seville, Spain | 287 |
| Nucleophilic displacement-reactions of the 4-sulfonyloxy group in derivatives having the D-manno configuration | |
| D. Cicero, O. Varela, and R. M. de Lederkremer (Buenos Aires, Argentina) | 295 |
| Analysis of a pyruvic acid acetal-containing polysaccharide by the reductive-cleavage method S. G. Zeller and G. R. Gray (Minneapolis, MN, U.S.A.) | 309 |
| Photolabeling of dextransucrase from <i>Streptococcus sanguis</i> with <i>p</i> -azidophenyl α-D-glucopyranoside | |
| S. F. Kobs and R. M. Mayer (Columbus, OH, U.S.A.) | 317 |
| Notes | |
| Enzymic synthesis of galactopyranosyl-L-serine derivatives using galactosidases D. Cantacuzene and S. Attal (Paris, France) | 327 |
| Synthesis of polymerizable anhydrodeoxyribose derivatives K. Hatanaka, Y. Yoshida, T. Yoshida, and T. Uryu (Tokyo, Japan) | 333 |
| Acceptor activity of affinity-immobilized dextransucrase from Streptococcus sanguis ATCC 10558 S. F. Kobs (Columbus, OH, U.S.A.) | 337 |
| Preliminary communication | |
| A facile total synthesis of ganglioside GM1b and its positional analog H. Prabhanjan, H. Ishida, M. Kiso, and A. Hasegawa (Gifu, Japan) | c1 |
| Author index | c7 |
| Subject index | с9 |
| Contents (Vol. 211) | c13 |

